

# TECH talk CE

THE NATIONAL CONTINUING EDUCATION PROGRAM FOR PHARMACY TECHNICIANS 1 CEU

FREE

ANSWER ONLINE FOR INSTANT RESULTS AT [WWW.CANADIANHEALTHCARENETWORK.CA](http://WWW.CANADIANHEALTHCARENETWORK.CA)

NOVEMBER 2011

## APPROVED FOR 1 CE UNIT



Approved for 1 CE unit by the Canadian Council on Continuing Education in Pharmacy. File # 1065-2011-265-I-T Tech. Not valid for CE credits after June 8, 2014.

Answer this CE online for instant results and accreditation. Visit [www.CanadianHealthcareNetwork.ca](http://www.CanadianHealthcareNetwork.ca)

### CE JUST FOR TECHNICIANS

Tech Talk CE is the only national continuing education program for Canadian pharmacy technicians.

As the role of the technician expands, use Tech Talk CE as a regular part of your learning portfolio. Note that a passing grade of 70% is required to earn the CE credit.

Tech Talk CE is generously sponsored by Teva. Download back issues at [www.CanadianHealthcareNetwork.ca](http://www.CanadianHealthcareNetwork.ca) or [www.tevacanada.com](http://www.tevacanada.com).

### ANSWERING OPTIONS

1. Answer the lesson online and get your results instantly at [www.CanadianHealthcareNetwork.ca](http://www.CanadianHealthcareNetwork.ca)
2. Use the reply card inserted with this CE lesson. Circle the answers on the card and mail in the prepaid, self-addressed card or fax to Mayra Ramos, 416-764-3937.

To pass this lesson, a grade of 70% (11 out of 15) is required. If you pass, you will receive 1 CEU. You will be advised of your results in a letter from *Tech Talk*. Please allow 8 to 12 weeks.

### CE FACULTY

#### CE Coordinator:

Margaret Woodruff, R.Ph., B.Sc.Pharm., MBA  
Humber College

#### Clinical Editor:

Lu-Ann Murdoch, B.Sc.Pharm.

#### Author:

Ema Ferreira, B.Pharm., M.Sc., Pharm.D., FSCPH

#### Reviewer:

Bobbi Thomas-Bailey, R.Ph.T., B.Ed.

# Treatment of Common Ailments During Pregnancy

by Ema Ferreira, B.Pharm., M.Sc., Pharm.D., FSCPH

## Learning Objectives:

Upon successful completion of this lesson, technicians will be able to:

1. Assist the pharmacist in identifying areas of counselling for prenatal care;
2. Identify women who are pregnant and may require the pharmacist's assistance; and
3. Describe the appropriate treatment for common disorders of pregnancy that may be treated with OTC medications.

### Introduction

Pharmacy technicians can play a valuable role in the care of pregnant women. By becoming familiar with optimal health habits, nutrition requirements and common ailments during pregnancy, they can help pharmacists provide preconception and prenatal care.

### Health habits

**Tobacco** Tobacco is associated with several adverse outcomes during pregnancy and ideally women should stop smoking before becoming pregnant.<sup>(1)</sup> In preconception/prenatal counselling, the pharmacy team should assess each woman's smoking status, explore her readiness to quit or cut down, identify strategies to help her, and refer her to the appropriate services for individual counselling.<sup>(2)</sup> As well, if appropriate, smoking cessation aids sold in pharmacies can be recommended or prescribed. Nicotine replacement products (e.g., transdermal patches and gums) can be used if deemed appropriate.<sup>(3)</sup>

**Alcohol use** Alcohol intake can cause birth defects, fetal and postnatal growth retardation, and central nervous system impairment.<sup>(2)</sup> Every woman who receives preconception counselling should be asked about alcohol consumption.

Since the exact amount of alcohol associated with fetal harm is not known, women who plan to become pregnant or are pregnant should abstain from alcohol.<sup>(2)</sup>

### Nutrition and vitamin supplements

Before pregnancy, women should be evaluated to ensure they have healthy eating habits, adequate intakes of calcium, vitamin D, folic acid and iron, and are not consuming excessive amounts of vitamin A (retinol; tolerable upper intake levels are 9,333-10,000 IU per day). Table 1 (available online) lists recommended daily allowances for calcium, folic acid, iron and vitamin D. Women planning a pregnancy should be encouraged to consume a healthy diet, according to Canada's Food Guide ([www.hc-sc.gc.ca/fn-an/food-guide-aliment/index\\_e.html](http://www.hc-sc.gc.ca/fn-an/food-guide-aliment/index_e.html)).<sup>(5)</sup>

Folic acid plays an important role in the prevention of congenital malformations, particularly neural tube defects (e.g., spina bifida). It may also help reduce other congenital anomalies, including cardiovascular, oral clefts, limb deformities and urinary malformations.<sup>(6)</sup> All women of child-bearing age should be counselled on the appropriate dose of folic acid to prevent congenital anomalies and the pharmacy technician can help to identify women who might benefit from this counselling.<sup>(7)</sup>

### Use of drugs during pregnancy

Pregnancy is normally 40 weeks in duration and is usually divided into three trimesters of 13 weeks each. The use of prescribed and over-the-counter (OTC) medications is common during pregnancy. In an American study, after excluding prenatal vitamins and iron supplements, 76.5% of pregnant women took at least one other medication, 62.8% used OTC medications, and 4.1% used herbal medications and/or alternative remedies. Multiple drug use occurred in 33.5% of patients, with up to 13.6% taking four or more remedies. In this study, the most common OTC medication taken by pregnant women was acetaminophen (39.6%), followed by antacids (32.9%) and ibuprofen (10.1%).<sup>(8)</sup>

### Treatment of common ailments

Pregnant women will consult the pharmacy team for different reasons (contraception, family planning, renewal of drugs, etc.), and pharmacists can provide them with useful advice and information. The pharmacy team should be able to recognize the signs and symptoms that require a medical referral and women should be advised to contact their physician at once if any of the signs listed in Table 2 appear. Such signs could indicate pregnancy loss, antepartum hemorrhage, preterm labour, and preterm rupture of membranes, hypertensive disorders of pregnancy, clots, infection, or hyperemesis gravidarum.

### Nausea and vomiting

The exact cause of nausea and vomiting during pregnancy is not known. The condition occurs in 75–80% of women, beginning early in the pregnancy and subsiding in a majority of women after the 14th week.<sup>(11)</sup> Even though it is often called morning sickness, it can occur at any time of the day. The recommendations in Table 3 may alleviate symptoms, although there is no evidence to prove the effectiveness of dietary changes.<sup>(5,10,11)</sup>

Medications can be used to control nausea and vomiting when the means mentioned in Table 3 do not work. A medical referral is required if symptoms affect daily activities, and medications can be prescribed to alleviate symptoms. OTC medications can be used to tide patients over until they see their physician.

Doxylamine/pyridoxine is the most widely used prescription antiemetic during

pregnancy<sup>(11)</sup>. This combination is efficacious and its safety during pregnancy is well-established. OTC antihistamines, such as dimenhydrinate, diphenhydramine and doxylamine, are generally regarded as safe during pregnancy.<sup>(12)</sup> Pyridoxine (vitamin B6) 10–25 mg three times a day may be effective to relieve nausea during pregnancy, with minimal side effects.<sup>(13)</sup>

Ginger is effective in the treatment of pregnancy-related nausea in doses ranging from 500–1500 mg per day.<sup>(13-15)</sup> Not all studies have evaluated pregnancy outcomes, although data suggest that ginger is safe during pregnancy, without causing a higher incidence of miscarriage or birth defects.<sup>(14,15)</sup> However, until all herbal products are well-regulated in Canada, ginger products should not be recommended routinely, since contamination with prescription drugs, heavy metals, and other substances is possible. If a patient insists on ginger, only products labelled with a Natural Product Number (NPN) should be recommended, since these have been evaluated by Health Canada for efficacy, safety and quality.

If a pregnant patient has severe symptoms and presents signs of dehydration (postural hypotension, dark urine, constant thirst, and dry skin), immediately refer her to the emergency room of the closest hospital.<sup>(5,11)</sup>

### Heartburn and acid reflux

More than 50% of pregnant women experience heartburn.<sup>(5,16,17)</sup> Symptoms usually appear during the first trimester and worsen as the pregnancy evolves.<sup>(16)</sup> Heartburn and indigestion are thought to be caused by the increase in hormones in early pregnancy, which slows digestion and relaxes the esophageal sphincter.<sup>(5,18)</sup> Increased intra-abdominal pressure due to the expansion of the uterus also contributes to the symptoms.<sup>(16)</sup> Simple dietary and lifestyle changes can relieve heartburn, such as eating frequent, small meals and eliminating causes of gastrointestinal discomfort (hot, spicy, fried, fatty foods, processed meats, caffeine, and alcohol). Also, women should avoid lying down immediately after eating, and rest and sleep with the head of the bed elevated (at least 15 cm or 6 inches).<sup>(19)</sup>

Despite dietary and lifestyle changes, some women will require pharmacological measures to relieve their heartburn.<sup>(16)</sup> Antacids containing aluminum hydroxide and

**TABLE 2 – Signs and symptoms that require a medical referral<sup>(9)</sup>**

- Bleeding from vagina, including spotting
- Fluid leaking from the vagina at any time before labour begins
- Sudden swelling or puffiness of the face, hands, or feet (particularly if it worsens or changes)
- Dizziness, lightheadedness, fainting spells, or loss of consciousness
- Headaches that are severe and last for a prolonged period of time
- Visual disturbances such as blurring, spots, flashes, or double vision
- Abdominal pain
- Chest pain
- Pain or burning when passing urine
- Chills, fever, or a rash following a fever
- Nausea or vomiting that lasts throughout the day
- Absence or decrease of fetal movement after the 24th week of pregnancy
- Signs of preterm labour (e.g., uterine contractions, vaginal leaking)
- Exposure to infectious diseases, including rubella, measles, hepatitis B, and sexually transmitted diseases including HIV
- Jaundice or dark-coloured urine

magnesium hydroxide, calcium carbonate or alginic acid at usual recommended doses can be used and are generally regarded as safe during pregnancy.<sup>(20)</sup>

Pregnant women should avoid products containing bismuth subsalicylate since they contain a salicylate, which is associated with fetal toxicity, including intrauterine growth retardation, premature closure of the ductus arteriosus in utero and pulmonary hypertension.<sup>(21)</sup>

Ranitidine (available over the counter) is the histamine-H2 blocker most widely studied in pregnancy, including during the first trimester, and it is not a major teratogen.<sup>(17,21,22)</sup> Less data are available on the use of other anti-H2 drugs.<sup>(21,23)</sup> Ranitidine at usual recommended doses reduces heartburn symptoms and may reduce the need for antacids.<sup>(24,25)</sup> Women whose symptoms are not relieved by antacids should consult their physician before taking ranitidine or other OTC anti-H2 drugs. Prescription proton pump inhibitors (PPIs), in particular omeprazole, can also be used if antacids and anti-H2 drugs are not sufficient.<sup>(26)</sup>

### Pain, headaches, backaches

Headaches can be caused by the increased nasal swelling and congestion (sinuses) related to pregnancy and can also be

**TABLE 3 – Tips to avoid or relieve nausea and vomiting of pregnancy<sup>(5)</sup>****Modification of eating habits**

- Try to eat a few crackers or a piece of bread before getting up in the morning.
- Eat appealing foods, in small quantities and frequently (every 2 to 3 hours).
- Eat lower-fat foods, such as poultry and fish, fruit, breads, cereals, rice, and potatoes and avoid fried foods.
- Avoid highly seasoned foods.
- To avoid hunger, do not skip meals.
- Drink fluids in small quantities and between meals to avoid fullness during meals.
- Have a snack before bedtime or during the night.

**Other tips**

- Get out of bed slowly—avoid sudden movements.
- Avoid strong smells and cooking odours.
- Brush teeth at other times of the day rather than immediately after meals.
- Stay well-rested and avoid fatigue.
- Avoid cigarette smoke.

induced by fatigue and stress.<sup>(5)</sup> Non-pharmacological methods to alleviate pain include relaxation, rest, eating regularly, and applying alternating hot and cold packs for sinus and tension headaches. Applying ice to the back of the neck can also help. Headaches associated with a fever, visual disturbances, or edema of the head and/or face should be referred to a physician.<sup>(5)</sup>

Backaches usually result from the pressure and weight of the enlarging uterus, the relaxation of ligaments (hormone-mediated), poor posture and/or excess weight.<sup>(5)</sup> Women should be advised to keep their weight gain within the recommended range and avoid shoes with poor support. Lifting and carrying objects correctly can be preventative and lessen symptoms. Pelvic tilt and dromedary droop exercises (bump or arch the back up like a cat while on hands and knees) are usually helpful. Massages also bring relief and relaxation.<sup>(5)</sup>

Acetaminophen is generally regarded as safe and is considered the pain reliever of choice during pregnancy.<sup>(21,27)</sup> Nonsteroidal anti-inflammatory drugs (NSAIDs), including ibuprofen and naproxen, have not been associated with congenital anomalies to date; occasional doses are generally regarded as safe during pregnancy before the third trimester.<sup>(21,27)</sup> After this period, NSAIDs, including ibuprofen and acetylsalicylic acid (ASA) at analgesic doses, can cause

oligohydramnios (decreased amniotic fluid) and pulmonary hypertension in the fetus, as well as other adverse effects.<sup>(27)</sup>

If pain persists after two or three days of adequate doses of acetaminophen or NSAIDs, women should be advised to consult their physician.

In a small study (doses were not specified), NSAID use was associated with an increased risk of miscarriage when taken around conception.<sup>(28)</sup> Therefore, avoiding NSAIDs around the time of conception might be advisable for patients who are planning a pregnancy.

**Constipation and hemorrhoids**

Constipation is related to progesterone changes during pregnancy, which reduce gastrointestinal motility, resulting in increased absorption of water from the stool and hardening of the stools.<sup>(29)</sup> Decreased physical activity during pregnancy and compression of the intestine by the enlarging uterus also worsen constipation.<sup>(29)</sup>

Avoiding refined foods as well as increasing the intake of high-fibre foods (e.g., whole-grain breads or cereals, bran cereals, fresh or dried fruit, nuts and vegetables) can be recommended.<sup>(5)</sup> Appropriate fluid intake and regular exercise can also improve symptoms.<sup>(5,29)</sup>

Laxatives can be used if non-pharmacological methods are insufficient. Bulk-forming laxatives (e.g., psyllium, calcium polycarboxylate or methylcellulose), at usual recommended doses, are generally regarded as safe during pregnancy. Sodium docusate and docusate calcium are emollient laxatives that are poorly absorbed and generally regarded as safe during pregnancy.<sup>(27,29)</sup> Recommended doses of lactulose, a hyperosmotic laxative, and stimulants such as sennosides, cascara sagrada, and bisacodyl can also be used when patients do not respond to initial treatment. Absorption of these laxatives is minimal and untoward pregnancy effects are unlikely.<sup>(27,29)</sup> Castor oil is contraindicated during pregnancy, as it can cause uterine contractions.<sup>(27,29)</sup> Mineral oil should also be avoided, since it can reduce lipid-soluble vitamin absorption.<sup>(29)</sup>

The pressure of the uterus on the pelvic veins and the straining that accompanies untreated constipation can cause hemorrhoids.<sup>(29)</sup> To avoid and relieve

hemorrhoids, patients should be counselled to prevent or treat constipation by following the above recommendations. Patients should avoid straining during bowel movements and use cold compresses to reduce hemorrhoid swelling, or warm sitz baths for comfort. Kegel exercises (pelvic floor exercises) may help to encourage venous return. Sleeping in the left-side lying position (not on the back) and avoidance of standing or sitting for prolonged periods also reduce pressure on the rectal veins. Use of topical medications or suppositories may be recommended. Hamamelis (also known as witch hazel) and glycerin compresses or zinc ointment (with or without pramoxine) are generally regarded as safe during pregnancy. If these measures are not useful or if rectal bleeding occurs, refer the patient to her physician.

Preparations containing shark liver oil should generally be avoided, since they contain vitamin A that can potentially be absorbed.<sup>(27,29)</sup> Topical hydrocortisone can be added to relieve itching or a rash; topical corticosteroids are deemed safe during pregnancy.<sup>(22,27)</sup>

**Vaginal infections**

Vaginal infections are common during pregnancy and women may seek advice on OTC antifungal treatments. Medical referral is important before recommending a treatment since some vaginal infections (e.g., bacterial vaginosis, *Trichomonas*) are associated with adverse pregnancy outcomes. In addition, first-lineazole antifungal treatments for vaginal candidiasis (*Candida albicans*) should be given for six to seven days; the shorter-course nonprescription products are not recommended during pregnancy.<sup>(30-31)</sup>

**The role of the pharmacy technician**

The pharmacy team has an important role to play in patient education in order to ensure the proper treatment of common ailments during pregnancy. First, women have to understand the importance of treatment to prevent complications. The risks of not treating a condition also have to be discussed. Secondly, women must receive information on drug use during pregnancy to maximize adherence. If a woman understands that a drug can be taken safely during pregnancy, she might be more willing to take it appropriately. The

information should be complete and unbiased. Pharmacists can ensure proper follow-up and recommend prophylaxis if necessary. They can also provide prenatal counselling regarding lifestyle modifications (healthy diet, exercise, tobacco, alcohol, drugs) and appropriate medication use during pregnancy. A collaborative approach to treating pregnant women is recommended. Pharmacy technicians can assist pharmacists in all these actions. In addition to the provision of medications and supportive counselling, pharmacy technicians can assist in several specific ways:

- Take a complete drug history including prescribed, OTC, natural products and recreational drugs;
- Identify patients who are planning to become pregnant or are pregnant;
- Support prevention of congenital anomalies by offering information on the use of folic acid;
- Ensure that the pharmacy carries an

assortment of prenatal vitamins containing the appropriate amounts of folic acid and other minerals and vitamins;

- Reinforce messages about the importance of adherence to prescribed therapy and lifestyle modifications, including smoking cessation;
- Monitor OTC sales and suggest that pregnant patients consult with the pharmacist for the most appropriate selection of nonprescription products;
- Encourage patients who have medical conditions to consult their doctor before becoming pregnant.

### Conclusion

The pharmacy team is very accessible and in a good position to counsel women on preconception and prenatal care. Pregnant women and women planning to be pregnant should be counselled about healthy eating habits and the risks of smoking, alcohol, and drug use. Healthy

nutrition is essential to achieve optimal fetal growth and maintain maternal health. According to current knowledge, women of childbearing age and pregnant women should pay special attention to the adequate intake of the essential nutrients iron, folic acid, and calcium. Nausea and vomiting, heartburn, headaches, back pain, constipation, hemorrhoids, and vaginal infections are common conditions that affect pregnant women. Non-pharmacological advice can help relieve symptoms; however, in many circumstances medications may be necessary. Several medications are generally regarded as safe during pregnancy and the pharmacy technician can assist the pharmacist to ensure drug adherence through proper counselling and information.

Table 1 and references are available at [www.CanadianHealthcareNetwork.ca](http://www.CanadianHealthcareNetwork.ca), CE section, Quick search CCCEP # 1065-2011-265-I-T

## QUESTIONS

### 1. Which statement is inaccurate?

- It is recommended that women who become pregnant abstain from alcohol since the exact amount associated with adverse pregnancy outcomes are unknown.
- Women planning to become pregnant should be encouraged to adapt healthy eating habits and should be informed about the benefits of adding a supplement of vitamin A to their daily intake.
- Tobacco smoking is associated with several negative pregnancy outcomes and if appropriate, smoking cessation aids should be recommended to pregnant women.
- Folic acid supplementation is recommended to all pregnant women in order to prevent congenital malformations.

### 2. What is the recommended daily dietary allowance of folic acid during pregnancy?

- 0.4 mg
- 0.6 mg
- 1 mg
- 5mg

Questions 3 and 4 refer to this case: Marissa is 25 weeks pregnant and she wants to have some advice to relieve her heartburn. She has no allergies and her pregnancy is going well. She smokes five cigarettes per day and takes prenatal vitamins daily.

### 3. What non-pharmacological advice would be appropriate to relieve her symptoms?

- Eat small meals
- Avoid eating before going to bed

- Avoid spicy foods
- Stop smoking
- All of the above

### 4. Which medication is not appropriate for the treatment of her acid reflux symptoms?

- Calcium carbonate tablets
- Magnesium and aluminum hydroxide liquid
- Bismuth subsalicylate liquid
- Alginic acid and aluminum hydroxide liquid

### 5. Which of following symptoms does not require a prompt medical referral?

- Spotting from the vagina
- Abdominal or chest pain
- Nausea before meals relieved by dimenhydrinate
- Dark-colored urine
- Headache not relieved by acetaminophen

Questions 6, 7 and 8 refer to the following case: A 32-year-old patient comes to your pharmacy to get some advice on what to do to relieve her nausea. She is seven weeks pregnant and has been nauseated for two weeks. She does not vomit but is bothered by the constant nausea. She will see her physician in one week.

### 6. What non-pharmacological advice can be given to her?

- Eat foods that are bland and tasteless.
- Skip meals to avoid fullness.
- Brush your teeth after each meal.
- Try to eat a piece of bread before getting up in the morning.

### 7. What pharmacological intervention would be best for the pharmacist to recommend to this woman at this point?

- Doxylamine 10 mg + pyridoxine 10 mg 2 tablets daily
- Ginger 500 mg po bid
- Dimenhydrinate 25 mg - 50 mg po q6h PRN
- Metoclopramide 10 mg po q6h PRN

The same patient returns to your pharmacy three weeks later after taking doxylamine 10 mg/vitamin B6 (pyridoxine) 10 mg tablets for two weeks (1 tablet morning and afternoon and 2 tablets at bedtime). She would like to renew her antiemetic prescription and talks to the pharmacy technician. She felt fine for the first two weeks but has started vomiting every day, feels dizzy when standing up and is constantly thirsty. You take her blood pressure and it is 90/50 mmHg.

### 8. What should the pharmacy technician do?

- Renew and fill the prescription.
- Inform the pharmacist of this situation, renew and fill her prescription.
- Sell her some ginger supplements to optimize her antiemetic treatment.
- Ask the woman to sit down in the waiting room to see the pharmacist and inform the pharmacist of the urgency of the situation.
- Recommend to the patient to take dimenhydrinate 50 mg po qid PRN with doxylamine 10 mg/vitamin B6 10 mg tablets.

## QUESTIONS (Continued)

9. A patient who is 30 weeks pregnant has a mild headache without any other symptoms and asks you if there is a drug that she can take to stop this headache. Her blood pressure is normal and she has no other symptoms. What would be the best choice for her?

- a) ibuprofen 400 mg po qid PRN
- b) acetaminophen 650 mg po q4h PRN
- c) acetylsalicylic acid 325- 650 mg po q4h PRN
- d) ibuprofen 200 mg po q6h PRN

Questions 10 and 11 refer to this case:

A pregnant patient of 28 weeks has been suffering from constipation for three weeks. She drinks plenty of fluids and eats fibre-rich foods but remains constipated. She wants your advice on laxatives and treatment for her hemorrhoids.

10. Which of the following laxatives is not recommended to treat constipation during pregnancy?

- a) sennosides
- b) docusate sodium
- c) psyllium
- d) mineral oil

11. To treat her hemorrhoids, the pharmacist should advise her to use:

- a) Shark liver oil containing ointment
- b) Pramoxine and zinc sulfate ointment
- c) Tea tree oil
- d) Hydrocortisone cream 0.5%

Please select the best answer for each question or answer online at [www.CanadianHealthcareNetwork.ca](http://www.CanadianHealthcareNetwork.ca) for instant results.

12. Bridget, 34 years old, is 23 weeks pregnant with her second child. Her first child was born prematurely at 32 weeks and is now a healthy two-year-old. She thinks she has a vaginal yeast infection. Which of the following statements is correct?

- a) Pregnant women rarely suffer from vaginal yeast infections.
- b) Symptoms of vaginal yeast infections are different during pregnancy.
- c) Other vaginal infections can be mistaken for a vaginal yeast infection, so she should be referred to her physician.
- d) The recommended treatment duration of a vaginal infection during pregnancy is usually shorter to minimize exposing the baby to antifungal drugs.

13. Caroline, 33 years old, is planning a pregnancy and consults you to get an appropriate vitamin supplement. She already takes the appropriate folic acid dose to prevent congenital anomalies. However, while talking to her, she tells you that she does not take any milk products or any calcium supplement. What amount of elemental calcium would she require to consume during pregnancy?

- a) At least 500 mg daily
- b) At least 1000 mg daily.
- c) At least 1500 mg daily
- d) She does not require a calcium supplement until the second half of her pregnancy

14. How can a pharmacy technician contribute to care for women planning a pregnancy?

- a) Take a complete drug history including prescribed, OTC and natural products
- b) Identify patients who are planning to become pregnant or are pregnant
- c) Ensure that the pharmacy carries an assortment of prenatal vitamins containing the appropriate amounts of folic acid and other minerals and vitamins
- d) Monitor OTC sales and suggest that women consult with the pharmacist for the most appropriate selection of nonprescription products
- e) All the above

15. A woman with hypertension is planning to become pregnant. A pharmacy technician can assist the pharmacist to ensure she takes appropriate actions to achieve a successful pregnancy. Which of the following is not appropriate?

- a) Encourage the woman to consult her doctor before becoming pregnant
- b) Reinforce messages about the importance of stopping medications before becoming pregnant
- c) Inform her of the availability of folic acid-containing vitamins to take before becoming pregnant
- d) Encourage a healthy lifestyle including smoking cessation if appropriate
- e) All of the above

## TECH talk CE

Presented by  
**pharmacy**  
practice

Sponsored by:  
**TEVA**

### Treatment of Common Ailments During Pregnancy

1 CEU • NOVEMBER 2011

CCCEP # 1065-2011-265-I-T Tech.

Not valid for CE credits after June 8, 2014.

### Now accredited by the Canadian Council on Continuing Education in Pharmacy

- |              |              |              |             |               |
|--------------|--------------|--------------|-------------|---------------|
| 1. a b c d   | 4. a b c d   | 7. a b c d   | 10. a b c d | 13. a b c d   |
| 2. a b c d   | 5. a b c d e | 8. a b c d e | 11. a b c d | 14. a b c d e |
| 3. a b c d e | 6. a b c d   | 9. a b c d   | 12. a b c d | 15. a b c d e |

First Name _____		Last Name _____	
Pharmacy Name _____			
Primary Licensing Province _____	Licence # _____	Secondary Licensing Province _____	Licence # _____
Home Address _____	City _____	Province _____	
Postal Code _____	Telephone _____	Fax _____	
Email _____		Year Graduated _____	

#### Type of practice

- Drug chain or franchise
- Grocery store pharmacy
- Banner
- Hospital pharmacy
- Independent
- Other (specify): \_\_\_\_\_
- Mass merchandiser

- Full-time technician
- Part-time technician

- Are you a certified technician?  
 Yes  No

Please help ensure this program continues to be useful to you by answering these questions.

- Do you now feel more informed about the treatment of common ailments during pregnancy?  Yes  No
- Was the information in this lesson relevant to you as a technician?  Yes  No
- Will you be able to incorporate the information from this lesson into your job as a technician?  Yes  No  N/A
- Was the information in this lesson...  Too basic  Appropriate  Too difficult
- How satisfied overall are you with this lesson?  
 Very  Somewhat  Not at all
- What topic would you like to see covered in a future issue? \_\_\_\_\_

**HOW TO ANSWER:** Answer ONLINE for immediate results at [www.CanadianHealthcareNetwork.ca](http://www.CanadianHealthcareNetwork.ca)

For information about CE marking, please contact Mayra Ramos at (416) 764-3879 or fax (416) 764-3937 or email [mayra.ramos@rci.rogers.com](mailto:mayra.ramos@rci.rogers.com). All other inquiries about Tech Talk CE should be directed to Karen Welds at (416) 764-3926 or [karen.welds@rci.rogers.com](mailto:karen.welds@rci.rogers.com).

**TABLE 1 - Recommended daily dietary allowances during pregnancy<sup>(6)</sup>**

Vitamin/mineral	Recommended daily dietary allowance
Calcium	1000-1300 mg*
Folic acid	0.6 mg
Iron	27 mg (elemental)
Vitamin D	600 IU

\* 1300 mg is recommended for women under the age of 18

**REFERENCES**

- DiFranza J, Aligne C, Weitzman M. Prenatal and postnatal environmental tobacco smoke exposure and children's health. *Pediatrics* 2004; 113(4):1007-1015.
- Kirkham C, Harris S, Grzybowski S. Evidence-based prenatal care: Part I. General prenatal care and counselling issues. *Am Fam Physician* 2005;71(7):1307-1560.
- Benowitz N, Dempsey D. Pharmacotherapy for smoking cessation during pregnancy. *Nicotine Tob Res* 2004;6 Suppl 2:S189-202.
- Wattendorf D, Muenke M. Fetal alcohol spectrum disorders. *Am Fam Physician* 2005;72(2):279-85.
- Health Canada. Nutrition for a healthy pregnancy: national guidelines for the childbearing years. Ottawa: Health Ministry; 1999.
- Health Canada. Dietary Reference Intakes. Available at: <http://www.hc-sc.gc.ca/fn-an/nutrition/reference/table/index-eng.php#rvv>. Accessed April 17, 2011.
- Wilson RD, Johnson JA, Wyatt P, et al. Pre-conceptional vitamin/folic acid supplementation 2007: the use of folic acid in combination with a multivitamin supplement for the prevention of neural tube defects and other congenital anomalies. *J Obstet Gynaecol Can* 2007;29(12):1003-26.
- Refuerzo J, Blackwell S, Sokol R, et al. Use of over-the-counter medications and herbal remedies in pregnancy. *Am J Perinatol* 2005;6(22):321-324.
- Health Canada. Care during pregnancy. In : Family-Centered Maternity and newborn care: national guidelines. Ottawa: Minister of Public Works and Government Services; 2000.
- Davis M. Nausea and vomiting of pregnancy: an evidence-based review. *J Perinat Neonatal Nurs* 2004;18(4):312-328.
- Arsenault MY, Lane CA. The management of nausea and vomiting during pregnancy. *J Gynaecol Obstet Can* 2002;24(10):817-823.
- Mazzotta P, Magee LA. Risk-benefit assessment of pharmacological and non-pharmacological treatments of nausea and vomiting of pregnancy. *Drugs* 2000;59(4):781-800.
- Jewell D, Young G. Interventions for nausea and vomiting in early pregnancy. *Cochrane Database Syst Rev* 2003(4):CD000145.
- Boone SA, Shields KM. Treating pregnancy-related nausea and vomiting with ginger. *Ann Pharmacother* 2005;39(10):1710-3.
- Smith C, Crowther C, Willson K, et al. A randomized controlled trial of ginger to treat nausea and vomiting in pregnancy. *Obstet Gynecol* 2004;103(4):639-45.
- Marrero JM, Goggin PM, de Caestecker JS, et al. Determinants of pregnancy heartburn. *Br J Obstet Gynaecol* 1992;99(9):731-4.
- Borum ML. Gastrointestinal diseases in women. *Med Clin North Am* 1998;82(1):21-50.
- Broussard CN, Richter JE. Treating gastro-oesophageal reflux disease during pregnancy and lactation: what are the safest therapy options? *Drug Saf* 1998;19(4):325-37.
- Katz PO, Castell DO. Gastroesophageal reflux disease during pregnancy. *Gastroenterol Clin North Am* 1998;27:153-167.
- Williamson C. Drugs in pregnancy. Gastrointestinal disease. *Best Pract Res Clin Obstet Gynaecol* 2001;15(6):937-52.
- Richter JE. Gastroesophageal reflux disease during pregnancy. *Gastroenterol Clin North Am* 2003;32(1):235-61.
- Briggs G, Freeman R, Yaffe SJ. *Drugs in Pregnancy and Lactation: A Reference Guide to Fetal and Neonatal Risk*. 8th ed. Philadelphia: Lippincott Williams & Wilkins, 2008.
- Conover EA. Herbal agents and over-the-counter medications in pregnancy. *Best Pract Res Clin Endocrinol Metab* 2003;17(2):237-51.
- Larson JD, Patatianian E, Miner PB Jr, et al. Double-blind, placebo-controlled study of ranitidine for gastroesophageal reflux symptoms during pregnancy. *Obstet Gynecol* 1997;90(1):83-7.
- Rayburn W, Liles E, Christensen H, et al. Antacids vs. antacids plus non-prescription ranitidine for heartburn during pregnancy. *Int J Gynaecol Obstet* 1999;66(1):35-7.
- Gill SK, O'Brien L, Einarson TR, et al. The safety of proton pump inhibitors (PPIs) in pregnancy: a meta-analysis. *Am J Gastroenterol* 2009;104:1541-5
- Ferreira E, ed. *Grossesse et allaitement: guide thérapeutique*. Montreal: Éditions Ste-Justine, 2007.
- Li DK, Liu L, Odouli R. Exposure to non-steroidal anti-inflammatory drugs during pregnancy and risk of miscarriage: population based cohort study. *BMJ* 2003;327(7411):368.
- Wald A. Constipation, diarrhea, and symptomatic hemorrhoids during pregnancy. *Gastroenterol Clin North Am* 2003;32(1):309-22, vii.
- Public Health Agency of Canada. Canadian guidelines on sexually transmitted infections 2008 (2010 update). [www.publichealth.gc.ca/sti](http://www.publichealth.gc.ca/sti) (accessed March 21, 2010).
- Center for Diseases Control and Prevention. Sexually transmitted diseases treatment guidelines, 2010. *MMWR* 2010;59(No RR12):56-61.