

PRODUCT MONOGRAPH
INCLUDING PATIENT MEDICATION INFORMATION

^{Pr} **NOVAMOXIN**

AMOXICILLIN CAPSULES, USP
Capsules, 250 mg and 500 mg amoxicillin (as amoxicillin trihydrate), Oral

AMOXICILLIN CHEWABLE TABLETS, USP
Tablets, 250 mg amoxicillin (as amoxicillin trihydrate), Oral

AMOXICILLIN GRANULES FOR ORAL SUSPENSION, USP
Suspension, 250 mg/5 ml amoxicillin (as amoxicillin trihydrate) after reconstitution, Oral

AMOXICILLIN SUGAR-REDUCED GRANULES FOR ORAL SUSPENSION, Teva Standard
Suspension, 250 mg/5 ml amoxicillin (as amoxicillin trihydrate) after reconstitution, Oral

Antibiotic

Teva Canada Limited
30 Novopharm Court
Toronto, ON
M1B 2K9
Canada
www.tevacanada.com

Date of Initial Authorization:
February 9, 1979

Date of Revision:
May 16, 2023

Submission Control Number: 268379

RECENT MAJOR LABEL CHANGES

3 SERIOUS WARNINGS AND PRECAUTIONS BOX	05/2023
7 WARNINGS AND PRECAUTIONS	05/2023

TABLE OF CONTENTS

PART I: HEALTH PROFESSIONAL INFORMATION	4
1 INDICATIONS	4
1.1 Pediatrics	4
1.2 Geriatrics.....	4
2 CONTRAINDICATIONS	4
3 SERIOUS WARNINGS AND PRECAUTIONS BOX	5
4 DOSAGE AND ADMINISTRATION	5
4.1 Dosing Considerations	5
4.2 Recommended Dose and Dosage Adjustment.....	5
4.3 Reconstitution	7
4.4 Administration	7
4.5 Missed Dose.....	7
5 OVERDOSAGE	7
6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING	8
7 WARNINGS AND PRECAUTIONS	9
7.1 Special Populations.....	12
7.1.1 Pregnant Women.....	12
7.1.3 Pediatrics	12
7.1.4 Geriatrics.....	12
8 ADVERSE REACTIONS	12
8.1 Adverse Reaction Overview.....	12
8.2 Clinical Trial Adverse Reactions.....	12
8.5 Post-Market Adverse Reactions	14
9 DRUG INTERACTIONS	14
9.4 Drug-Drug Interactions.....	14
9.5 Drug-Food Interactions.....	14
9.6 Drug-Herb Interactions.....	15
9.7 Drug-Laboratory Test Interactions	15
10 ACTION AND CLINICAL PHARMACOLOGY	15
10.1 Mechanism of Action.....	15
10.3 Pharmacokinetics	15
11 STORAGE, STABILITY AND DISPOSAL	16
PART II: SCIENTIFIC INFORMATION	17
13 PHARMACEUTICAL INFORMATION	17
14 CLINICAL TRIALS	17
14.2 Comparative Bioavailability Studies - Tablets	17
15 MICROBIOLOGY	18
16 NON-CLINICAL TOXICOLOGY	19

PATIENT MEDICATION INFORMATION..... 23

PART I: HEALTH PROFESSIONAL INFORMATION

1 INDICATIONS

Novamoxin (amoxicillin) may be indicated in the treatment of infections due to susceptible strains of the following micro-organisms: Gram-negative organisms: *H. influenzae*, *P. mirabilis* and *N. gonorrhoeae*. Gram-positive organisms: Streptococci (including *Streptococcus faecalis* and *Streptococcus pneumoniae*).

Amoxicillin is not active against *Pseudomonas aeruginosa*, indole-positive *Proteus* species, *Serratia marcescens*, *Klebsiella* and *Enterobacter* species.

In emergency cases, where the causative organism is not yet identified, therapy may be initiated with amoxicillin on the basis of clinical judgment while awaiting bacteriologic tests to determine its antimicrobial sensitivity.

Novamoxin may be indicated as a prophylaxis against alpha-hemolytic (Viridan's group) Streptococci before dental, oral or upper respiratory tract surgery or instrumentation.

It may be also indicated as a prophylaxis of bacterial endocarditis in patients with any of the following conditions: congenital cardiac malformations, rheumatic and other acquired valvular lesions, prosthetic heart valves, previous history of bacterial endocarditis, hypertrophic cardiomyopathy, surgically constructed systemic pulmonary shunts, mitral valve prolapse with valvular regurgitation or mitral valve prolapse without valvular regurgitation but associated with thickening and/or redundancy of the valve leaflets.

To reduce the development of drug-resistant bacteria and maintain the effectiveness of Novamoxin and other antibacterial drugs, Novamoxin should be used only to treat infections that are proven or strongly suspected to be caused by susceptible bacteria. When culture and susceptibility information are available, they should be considered in selecting or modifying antibacterial therapy. In the absence of such data, local epidemiology and susceptibility patterns may contribute to the empiric selection of therapy.

1.1 Pediatrics

Pediatrics (<18): (See [4.2 Recommended Dose and Dosage Adjustment](#), [7.1.3 Pediatrics](#)).

1.2 Geriatrics

Geriatrics: (See [7.1.4 Geriatrics](#))

2 CONTRAINDICATIONS

NOVAMOXIN is contraindicated in:

- patients who are hypersensitive to this drug or to any ingredient in the formulation or component of the container.
- patients with a history of a previous hypersensitivity reaction to any of the penicillins or cephalosporins.
- in cases where infectious mononucleosis is either suspected or confirmed.

3 SERIOUS WARNINGS AND PRECAUTIONS BOX

Serious Warnings And Precautions Box

- Hypersensitivity: Serious and occasionally fatal hypersensitivity (anaphylactic) and severe cutaneous adverse reactions (SCAR) have been reported in patients receiving therapy with beta-lactams, including amoxicillin. (See [7 WARNINGS AND PRECAUTIONS, Hypersensitivity](#) and [7 WARNINGS AND PRECAUTIONS, Skin](#)).

4 DOSAGE AND ADMINISTRATION

4.1 Dosing Considerations

Periodic assessment of renal, hepatic and hematopoietic functions should be made during prolonged therapy with Novamoxin (amoxicillin).

4.2 Recommended Dose and Dosage Adjustment

Usual Dosage

Infections of the upper respiratory tract (ear, nose and throat) due to susceptible strains of streptococci (beta-hemolytic and *Streptococcus pneumoniae*), non penicillinase-producing staphylococci and *H. influenzae*.

Infections of the urinary tract due to *Proteus mirabilis* and *Streptococcus faecalis*.

Infections of the skin and soft-tissues due to streptococci and staphylococci (non penicillinase producing).

Adults: 250 mg every 8 hours

Children <20 kg: 20 mg/kg/day in divided doses every 8 hours. This dosage should not exceed the recommended adult dosage.

Children weighing 20 kg or more should be dosed according to the adult recommendations.

In severe infections or infections associated with organisms where sensitivity determinations require higher blood concentrations: 500 mg every 8 hours for adults, and 40 mg/kg/day in divided doses every 8 hours for children less than 20 kg may be needed.

Infections of the lower respiratory tract, due to susceptible strains of the causative organism and acute otitis media.

Adults: 500 mg every 8 hours
Children <20 kg: 40 mg/kg/day in divided doses every 8 hours. This dosage should not exceed the recommended adult dosage.

Children weighing 20 kg or more should be dosed according to the adult recommendations.

Urethritis due to nonpenicillinase producing *N. gonorrhoeae* acquired in area with active monitoring for resistance to penicillin and where the percentage of penicillin-resistant isolates is <3.0%:

Adults and children >45 kg: (3 g as a single oral dose); 1 g of oral probenecid should be administered concomitantly as well as appropriate therapy for presumptive or proven infection with *C. trachomatis*.

Children <45 kg: a single 50 mg/kg dose (maximum 3 g) given with a single 25 mg/kg (up to 1 g) dose of probenecid. However, probenecid is not recommended in children under 2 years of age. Appropriate therapy of presumptive or proven infection with *C. trachomatis* should be included as well. Cases of gonorrhoea with a suspected lesion of syphilis should have darkfield examinations before receiving amoxicillin, and monthly serological tests for a minimum of four months.

For prevention of endocarditis:

Adults: 3 g orally 1 hour before procedure; then 1.5 g 6 hours after the initial dose.

Children: 50 mg/kg (not to exceed adult dose) orally 1 hour before procedure; then 25 mg/kg 6 hours after the initial dose.

It should be recognized that in the treatment of chronic urinary tract infections, frequent bacteriological and clinical appraisals are necessary. Smaller doses than those recommended above should not be used. Even higher doses may be needed at times and in stubborn infections therapy may be required for several weeks. It may be necessary to continue clinical and/or bacteriological follow-up for several months after cessation of therapy. Except for gonorrhoea, treatment should be continued for a minimum of 48 to 72 hours beyond the time that the patient becomes asymptomatic or evidence of bacterial eradication has been obtained. It is recommended that there be at least 10 days treatment for any infection caused by beta-

hemolytic streptococci to prevent the occurrence of acute rheumatic fever or glomerulonephritis.

Dosage Adjustment

Renal Impairment: Because amoxicillin is excreted mostly by the kidney, the dosage for patients with renal impairment should be reduced in proportion to the degree of loss of renal function.

4.3 Reconstitution

Directions for Dispensing Oral Suspension:

Prepare these formulations at the time of dispensing. For ease in preparation, add water to the bottle in two portions and shake well after each addition. Add the total amount of water as directed on the labeling of the package being dispensed.

The reconstituted formulation is stable for 14 days under refrigeration (6°C) or 7 days at room temperature (25°C).

Novamoxin Granules for Oral Suspension 250 mg/5 mL: After reconstitution each 5 mL suspension contains amoxicillin trihydrate equivalent to 250 mg amoxicillin. Sugar content/5 mL: 250 mg suspension 2.75 g equivalent to 11 .0 cal.

Novamoxin Sugar-Reduced Granules for Oral Suspension 250 mg/5 mL: Each 5 mL of reconstituted suspension contains amoxicillin trihydrate equivalent to 250 mg amoxicillin.

4.4 Administration

In order to obtain optimal absorption of drug from Novamoxin capsules they should be administered between meals with a glass of water (250 mL or 8 fl. oz.).

4.5 Missed Dose

Patients should be instructed to take NOVAMOXIN at the next scheduled dose and not take two doses at the same time if they miss a dose.

5 OVERDOSAGE

Treatment of overdosage would likely be needed only in patients with severely impaired renal function, since patients with normal kidneys excrete penicillins at a fast rate. Hemodialysis would, therefore, represent the main form of treatment.

Activated charcoal may be administered to aid in the removal of unabsorbed drug. General supportive measures are recommended.

For management of a suspected drug overdose, contact your regional poison control centre.

6 DOSAGE FORMS, STRENGTHS, COMPOSITION AND PACKAGING

Table-1 – Dosage Forms, Strengths, Composition and Packaging

Route of Administration	Dosage Form / Strength / Composition	Non-medicinal Ingredients
Oral	Capsules, 250 mg and 500 mg	Colloidal Silicon Dioxide, Dry-Flo Starch, Magnesium stearate, Sodium lauryl sulfate. Gelatine Capsules (EGC Size #2) and (EGC Size #0) : D&C Yellow # 10, FD & C Blue #1, FD & C Red #3, FD & C Red #40, FD & C Yellow # 6, Titanium Dioxide, Gelatin and White S-1-7085 (250 mg) / White SB-0007P (500 mg)
Oral	Tablets, 250 mg	Artificial Cherry Flavor, Aspartame, Colloidal Silicon Dioxide, Dipac White (Compressible Sugar), D&C Red #30 AL Lake, FD&C Red #40 Lake, Magnesium Stearate, Mannitol, Microcrystalline Cellulose
Oral	Granules for Oral Suspension, 250 mg/5 mL	Artificial Cherry Raspberry Flavor, FD&C Red #40, Silicon Dioxide, Sodium Benzoate, Sodium Citrate, Sucrose, Xanthan Gum
Oral	Sugar-Reduced Granules for Oral Suspension, 250 mg/5 mL	Colloidal Silicon Dioxide, Mannitol 2080, Nipagin-M Sodium, Nipazol-M Sodium, Sodium Citrate, Sucrose, Xanthan Gum For 250 mg/5 mL Sugar-Reduced Granules for Oral Suspension: D&C Yellow #10 & Artificial Banana Flavour

Hard Gelatin Capsules:

NOVAMOXIN 250 mg Capsules: 250 mg amoxicillin (as the trihydrate) in # 2 capsules with opaque scarlet cap and yellow body. Printed white NOVO and 250 on opposing cap and body portions.

NOVAMOXIN 500 mg Capsules: 500 mg amoxicillin (as the trihydrate) in #0 capsules with opaque scarlet cap and yellow body. Printed white NOVO and 500 on opposing cap and body portions.

Bottles of 100 capsules for 250 mg
Bottles of 100 and 500 capsules for 500 mg

Novamoxin Granules for Oral Suspension 250 mg/5 mL:

Each 5 mL of reconstituted suspension contains amoxicillin trihydrate equivalent to 250 mg amoxicillin.

250 mg supplied in bottles of 100 mL and 150 mL.

Novamoxin Sugar-Reduced Granules for Oral Suspension 250 mg/5 mL:

Each 5 mL of reconstituted suspension contains amoxicillin trihydrate equivalent to 250 mg amoxicillin.

250 mg supplied in bottles of 100 mL and 150 mL.

Chewable Tablets:

NOVAMOXIN 250 mg is available as a rose coloured, mottled, oval shaped, single-scored compressed tablet with cherry flavour, containing 250 mg of amoxicillin as the trihydrate. Engraved 'NOVO' with scoreline on one side and '250' on the other side.

Supplied: 100 and 500 for 250 mg.

7 WARNINGS AND PRECAUTIONS

General

If superinfections with mycotic or bacterial pathogens occur (usually involving *Aerobacter*, *Pseudomonas* or *Candida*) treatment with Novamoxin should be discontinued and appropriate therapy instituted.

Novamoxin is contraindicated in cases where infectious mononucleosis is either suspected or confirmed (see [2 CONTRAINDICATIONS](#)). A morbilliform rash following the use of ampicillin in patients with infectious mononucleosis has been well documented and has also been reported to occur following the use of amoxicillin.

Cardiovascular

Kounis syndrome, a serious allergic reaction that can result in myocardial infarction, can occur as chest pain in association with an allergic reaction to amoxicillin.

Gastrointestinal

Clostridium difficile-associated disease

Clostridium difficile-associated disease (CDAD) has been reported with use of many antibacterial agents, including amoxicillin (see [8 ADVERSE REACTIONS](#)). CDAD may range in severity from mild diarrhea to fatal colitis. It is important to consider this diagnosis in patients who present with diarrhea, or symptoms of colitis, pseudomembranous colitis, toxic megacolon, or perforation of colon subsequent to the administration of any antibacterial agent. CDAD has been reported to occur over 2 months after the administration of antibacterial agents.

Treatment with antibacterial agents may alter the normal flora of the colon and may permit overgrowth of Clostridium difficile. C. difficile produces toxins A and B, which contribute to the development of CDAD. CDAD may cause significant morbidity and mortality. CDAD can be refractory to antimicrobial therapy.

If the diagnosis of CDAD is suspected or confirmed, appropriate therapeutic measures should be initiated. Mild cases of CDAD usually respond to discontinuation of antibacterial agents not directed against Clostridium difficile. In moderate to severe cases, consideration should be given to management with fluids and electrolytes, protein supplementation, and treatment with an antibacterial agent clinically effective against Clostridium difficile. Surgical evaluation should be instituted as clinically indicated, as surgical intervention may be required in certain severe cases.

Hematologic

Periodic assessment of hematopoietic function should be made during prolonged therapy with Novamoxin (amoxicillin).

Hepatic/Biliary/Pancreatic

A moderate rise in serum glutamic oxaloacetic transaminase (SGOT) has been noted. Periodic assessment of hepatic function should be made during prolonged therapy with Novamoxin (amoxicillin). (See [7.1.3 Pediatrics](#); [8.2 Clinical Trial Adverse Reactions](#))

Susceptibility/Resistance

Development of Drug Resistant Bacteria

Prescribing Novamoxin in the absence of a proven or strongly suspected bacterial infection is unlikely to provide benefit to the patient and risks the development of drug-resistant bacteria.

Immune

Hypersensitivity

Serious and occasionally fatal hypersensitivity (anaphylactoid) reactions have been reported in patients on penicillin therapy. Although anaphylaxis is more frequent following parenteral therapy, it has occurred in patients following oral dosing of penicillins. These reactions are more likely to occur in patients with a history of hypersensitivity to beta-lactams and individuals with a history of sensitivity to multiple allergens. There have been well-documented reports of individuals with a history of penicillin hypersensitivity reactions who have experienced severe hypersensitivity reactions when treated with cephalosporins. Before initiating therapy with a penicillin, careful inquiry should be made concerning previous hypersensitivity reactions to penicillins, cephalosporins and other allergens. If an allergic reaction occurs, administration of Novamoxin (amoxicillin) should be discontinued and appropriate therapy instituted.

Serious anaphylactoid reactions require immediate emergency treatment with epinephrine. Oxygen, intravenous steroids, and airway management, including intubation, should also be administered as indicated.

Monitoring and Laboratory Tests

Periodic assessment of renal, hepatic and hematopoietic functions should be made during prolonged therapy with Novamoxin (amoxicillin).

Abnormal prolongation of prothrombin time (increased international normalized ratio (INR)) has been reported in patients receiving amoxicillin and oral anticoagulants. Appropriate monitoring should be undertaken when amoxicillin and oral anticoagulants are prescribed concurrently, particularly upon initiation or cessation of concurrent administration. Adjustments in the dose of oral anticoagulants may be necessary to maintain the desired level of anticoagulation.

Renal

Because amoxicillin is excreted mostly by the kidney, the dosage for patients with renal impairment should be reduced in proportion to the degree of loss of renal function

Periodic assessment of renal function should be made during prolonged therapy with Novamoxin (amoxicillin).

Skin

Severe Cutaneous Adverse Reactions

Severe cutaneous adverse reactions (SCAR) such as acute generalized exanthematous pustulosis (AGEP), drug reaction with eosinophilia and systemic symptoms (DRESS), Stevens-Johnson

syndrome (SJS), and toxic epidermal necrolysis (TEN) have been reported in association with beta-lactam treatment. When SCAR is suspected, NOVAMOXIN should be discontinued and appropriate therapy and/or measures should be taken.

7.1 Special Populations

7.1.1 Pregnant Women

The safety of Novamoxin in the treatment of infections during pregnancy has not been established. If the administration of Novamoxin to pregnant patients is considered to be necessary, its use requires that the potential benefits be weighed against the possible hazards to the fetus.

7.1.3 Pediatrics

Pediatrics (<18): A moderate rise in serum glutamic oxaloacetic transaminase (SGOT) has been noted, particularly in infants, but the significance of this finding is not known.

7.1.4 Geriatrics

Use in the Elderly: There are no known specific precautions for the use of amoxicillin in the elderly.

Amoxicillin is known to be substantially excreted by the kidney, and the risk of toxic reactions to this Novamoxin may be greater in patients with impaired renal function. Because elderly patients are more likely to have decreased renal function, care should be taken in dose selection, and it may be useful to monitor renal function.

8 ADVERSE REACTIONS

8.1 Adverse Reaction Overview

As with other penicillins, it may be expected that untoward reactions will be related to sensitivity phenomena. They are more likely to occur in individuals who have previously demonstrated hypersensitivity to penicillins and cephalosporins and in those with a history of allergy, asthma, hay fever or urticaria.

8.2 Clinical Trial Adverse Reactions

The following adverse reactions have been reported as associated with the use of Novamoxin:

Gastrointestinal - Nausea, vomiting and diarrhea, hemorrhagic and pseudomembranous colitis. Clostridium difficile-associated disease (CDAD) has been reported with use of many antibacterial

agents, including amoxicillin. Glossitis, black "hairy" tongue and stomatitis, mucocutaneous candidiasis, tooth discoloration (brown, yellow or gray staining); most reports occurred in pediatric patients. Discoloration was reduced or eliminated with brushing or dental cleaning in most cases.

Hypersensitivity Reactions - Skin rashes have been reported frequently. Less commonly, a few cases of serum sickness like reactions including urticaria, erythema, erythema multiforme, angioneurotic edema, pruritus have been reported. Rarely, Stevens-Johnson syndrome, toxic epidermal necrolysis, bullous dermatitis, exfoliative dermatitis, acute generalized exanthematous pustulosis, hypersensitivity vasculitis have been reported.

Anaphylaxis is the most serious reaction experienced and has usually been associated with the parenteral dosage form.

NOTE: Urticaria, other skin rashes, and serum sickness-like reactions may be controlled with antihistamines and if necessary, systemic corticosteroids. Whenever such reactions occur, Novamoxin (amoxicillin) should be discontinued unless, in the opinion of the physician, the condition being treated is life threatening and amenable only to amoxicillin therapy. Serious anaphylactic reactions require the immediate use of epinephrine, oxygen and intravenous steroids.

Hepatobiliary - A moderate rise in serum glutamic oxaloacetic transaminase (SGOT) has been noted, particularly in infants, but the significance of this finding is not known. Transient increases in serum alkaline phosphatase and lactic dehydrogenase levels have also been observed but they returned to normal on discontinuation of amoxicillin. Reports have also been seen of hepatic dysfunction including cholestatic jaundice, hepatic cholestasis, acute cytolytic hepatitis,

Hemic and Lymphatic Systems - Anemia thrombocytopenia, thrombocytopenic purpura, eosinophilia, leukopenia, neutropenia and agranulocytosis have been reported during therapy with the penicillins. These reactions are usually reversible on discontinuation of therapy and are believed to be a hypersensitivity phenomena. Reports have also been seen of anemia including hemolytic anemia.

Central Nervous System - As with other penicillins, acute and chronic toxicity is not a clinical problem. Although penicillins do not normally cross the blood-brain barrier to any substantial extent, if massive doses are given (several grams per day) to elderly patients, patients with inflamed meninges or patients with impaired renal function, toxic reactions are likely to occur. At extremely high doses, convulsions can occur. When penicillin reaches a high concentration in the cerebrospinal fluid, neurotoxic symptoms consisting of myoclonia, convulsive seizures and depressed consciousness may occur. Unless administration of the drug is stopped or its dosage reduced, the syndrome may progress to coma and death. Dizziness, hyperkinesias, hyperactivity, agitation, anxiety, insomnia, confusion, and behavioural changes have also been reported.

Skin and Appendages - erythematous maculopapular rash.

Renal - Crystalluria. Interstitial nephritis (oliguria, proteinuria, hematuria, hyaline casts, pyuria) and nephropathy are infrequent and usually associated with high doses of parenteral penicillins; however, this has occurred with all of the penicillins. Such reactions are hypersensitivity responses and are usually associated with fever, skin rash and eosinophilia. Elevations of creatinine or blood urea nitrogen may occur.

8.5 Post-Market Adverse Reactions

Central Nervous System: Amoxicillin can lead to cases of aseptic meningitis of unknown frequency.

Other immune system disorders: Kounis syndrome

9 DRUG INTERACTIONS

9.4 Drug-Drug Interactions

Methotrexate: Penicillins compete with renal tubular secretion of methotrexate, resulting in decreased clearance of methotrexate. Concomitant use may increase methotrexate serum concentrations, with increased risk of toxicity.

Probenecid: Probenecid inhibits the renal tubular excretion of amoxicillin. Concurrent use of amoxicillin and probenecid may result in increased and prolonged blood levels of amoxicillin.

Warfarin: Abnormal prolongation of prothrombin time (increased international normalized ratio [INR]) has been reported in patients receiving amoxicillin and warfarin. Appropriate monitoring should be undertaken when warfarin is prescribed concurrently. Adjustments in the dose of oral anticoagulants may be necessary to maintain the desired level of anticoagulation.

Oral Contraceptives: Novamoxin may affect the gut flora, leading to lower estrogen reabsorption and reduced efficacy of combined oral estrogen/progesterone contraceptives.

Tetracyclines: Bacteriostatic action of tetracyclines may inhibit bactericidal activity of penicillins.

9.5 Drug-Food Interactions

Amoxicillin is stable in the presence of gastric acid. Novamoxin is rapidly and well absorbed after oral administration to fasting subjects. It was found in a recent study that peak serum antibiotic levels were reduced by 50% in subjects receiving amoxicillin immediately following a standard meal. Reducing the dose-water volume given with amoxicillin from 250 to 25 mL in fasted subjects also caused a significant reduction in serum amoxicillin levels. This may be due to the

low water solubility of amoxicillin trihydrate (1 g in 370 mL water). In addition, food ingestion immediately before dosing also reduced the urinary excretion.

9.6 Drug-Herb Interactions

Interactions with herbal products have not been established.

9.7 Drug-Laboratory Test Interactions

Novamoxin may:

- cause false-positive reactions when testing for the presence of glucose in urine.
- distort assay results for estriol in pregnant women.

10 ACTION AND CLINICAL PHARMACOLOGY

10.1 Mechanism of Action

Novamoxin (amoxicillin) exerts its bactericidal action by interfering with bacterial cell wall synthesis.

10.3 Pharmacokinetics

Peak serum levels are attained between 1 and 2 hours after drug administration. Amoxicillin diffuses readily into most body tissues and fluids, with the exception of brain and spinal fluid. Amoxicillin is excreted largely unchanged in the urine while 10-25% of the administered dose is excreted in the form of penicilloic acid. The excretion of amoxicillin can be delayed by concurrent administration of probenecid. Amoxicillin is not highly protein bound. In blood serum, amoxicillin is approximately 17-18% protein bound compared to 59% for penicillin G.

The following amoxicillin mean serum levels were found following the administration of 250 mg capsules of Novamoxin to 12 healthy adult volunteers:

Time (hr.)	0.5	1.0	1.5	2	3	4	5	7
Mean Serum Levels ($\mu\text{g/mL}$)	0.81	2.96	3.17	3.10	2.22	1.12	0.50	0.11

Peak blood serum levels averaged $3.8\mu\text{g/mL}$ (range 2.35 to 6.38) and the T_{max} was 1.50 hr. The mean biological half-life ($t_{1/2}$) was found to be 55.8 minutes with a mean elimination rate constant K_{el} of 0.7456 hr.^{-1} .

Twelve normal male subjects participated in a bioavailability study of Novamoxin Granules for Suspension. Each subject was given 5 mL (250 mg) of reconstituted Novamoxin Granules for Suspension in a single dose.

The following amoxicillin mean serum levels were found:

Time (hr.)	0.5	1.0	1.5	2	3	4	5	7
Mean Serum Levels ($\mu\text{g/mL}$)	3.26	4.19	3.40	2.56	1.65	0.98	0.43	0.10

Peak plasma concentrations from 2.65 to 5.75 $\mu\text{g/mL}$ were obtained with a mean C_{max} of $4.24 \pm 0.74 \mu\text{g/mL}$. The time required to reach peak concentrations ranged from 0.5 to 1.5 hours, with a T_{max} mean of $1.00 \pm 0.21 \text{ hr}$.

The AUC's calculated for 0 to 7 hours ranged from 8.475 to 12.865 $\mu\text{g-hours/mL}$. The mean AUC was $10.713 \pm 1.443 \mu\text{g-hours/mL}$. The mean biological half-life for Novamoxin Granules for Suspension was 26.4 minutes. The mean elimination rate constant (K_{el}) was 1.57 hour^{-1} .

The administration of 500 mg amoxicillin to healthy fasting subjects has been reported to produce peak mean serum levels of 10.8 $\mu\text{g/mL}$ and 6.75 $\mu\text{g/mL}$. Additional studies in healthy volunteers with normal renal function receiving 500 mg doses, indicated that peak serum levels could vary from 5.0 to 10.8 $\mu\text{g/mL}$. Serum amoxicillin half-life values reported in the literature vary from 1-1.3 hours. About 60-80% of an oral dose of amoxicillin is excreted in the urine. In the presence of renal impairment the serum half-life increases (between 7 and 10 hours), necessitating a reduction in the dosage administered.

11 STORAGE, STABILITY AND DISPOSAL

Capsules and Tablets: Store between 15°C and 30°C. Unit dose strips should be stored between 15°C and 25°C and protected from high humidity.

Granules for Oral Suspension: Store at room temperature (between 15°C -30°C). Keep bottle tightly closed.

Reconstituted Solution: The reconstituted formulation is stable for 14 days under refrigeration (between 2°C -8°C) or 7 days at room temperature (between 15°C -30°C).

PART II: SCIENTIFIC INFORMATION

13 PHARMACEUTICAL INFORMATION


DRUG SUBSTANCE:

Trade Name: NOVAMOXIN

Proper Name: Amoxicillin Trihydrate

Chemical Name: Trihydrate of 6-[D-(-)-alpha-amino-4-hydroxyphenyl-acetamido] penicillanic acid.

Structural Formula:



Molecular Formula and molecular mass: $C_{16}H_{19}N_3O_5S \cdot 3H_2O$, 419.5

Description: Amoxicillin trihydrate is a white practically odourless crystalline powder, slightly soluble in water and in methanol; insoluble in benzenes, in chloroform and in ether.

14 CLINICAL TRIALS

14.2 Comparative Bioavailability Studies - Tablets

A comparative two-way, single-dose bioavailability study was carried out in sixteen healthy, male volunteers on two 250 mg amoxicillin tablet products, Novamoxin 250 mg and Amoxil 250 mg chewable tablets. The pharmacokinetic data of fifteen subjects calculated for the Novamoxin and Amoxil chewable tablet formulations is tabulated below:

Amoxicillin Trihydrate Chewable Tablets 250 mg				
From measured data				
Geometric Mean				
Arithmetic Mean (C.V.%)				
Parameter	Test*	Reference†	% Ratio of Geometric Means	95% Confidence Interval [§]
AUC _T (µg*h/mL)	20.3 20.4 (10)	20.5 20.7 (16)	99	91.8 – 108.4
AUC _I (µg• h/mL)	20.5 20.5 (10)	20.5 20.8 (16)	100	91.8 – 108.5
C _{max} (µg/mL)	7.4 7.5 (13)	7.7 8.0 (26)	96	84.8 – 109.1
T _{max} [§] (h)	1.37 (0.40)	1.43 (0.46)		-
T _{1/2} [§] (h)	0.95 (0.11)	0.92 (0.09)		-

*Novamoxin 250 mg Chewable Tablet (Novopharm Ltd.).

†Amoxil® 250 mg Chewable Tablet (purchased in Canada).

§Expressed as the arithmetic mean (standard deviation) only.

15 MICROBIOLOGY

In vitro studies with amoxicillin have demonstrated the susceptibility of the following gram-positive bacteria: beta-hemolytic streptococci, *Streptococcus pneumoniae*, *D. pneumoniae*, non-penicillinase-producing staphylococci, and *Streptococcus faecalis*. It is active *in vitro* against many strains of *Haemophilus influenzae*, *Neisseria gonorrhoeae* and *Proteus mirabilis*. Because amoxicillin does not resist destruction by penicillinase, it is not effective against penicillinase-producing bacteria, particularly resistant staphylococci.

Amoxicillin is not active against all *Pseudomonas aeruginosa*, indole-positive *Proteus* species, *Serratia marcescens*, *Klebsiella*, and *Enterobacter* species.

Disc Susceptibility Tests: Quantitative methods that involve the measurement of the diameters of zones of inhibition can be used to estimate micro-organism sensitivity to a particular antibiotic. A procedure which involves the use of discs impregnated with a particular antibiotic has been described for the ampicillin class of antibiotics. Interpretations correlate diameters of the zones of inhibition with MIC values for amoxicillin. With this procedure, using a 10 µg disc, a zone of 29 mm or more is classified as "susceptible" and indicates that the infecting organism is likely to respond to therapy. A zone of 20 mm or less is classified as "resistant" and indicates that the infecting organism is not likely to respond to therapy. A zone of 21 -28 mm is classified

as "intermediate susceptibility" and indicates that the organism would be susceptible if high dosages are used, or if the infection is confined to tissues and fluids (e.g., urine), in which antibiotic levels are attained.

The *in vitro* activity of amoxicillin against selected organisms has been reported by Sutherland *et al.* and Sabto *et al.* shown in the following tables:

Table I. *In Vitro* Activity of Amoxicillin Against Gram-Positive Cocci, H. Influenzae and N. Gonorrhoeae

Organism	No. of Strains	Minimum Inhibitory Concentration ($\mu\text{g}/\text{mL}$)								
		.005	0.01	0.02	0.03	0.05	0.12	0.25	0.5	1.0
Staphylococcus aureus	29					3	20	6		
Beta-hemolytic streptococci	28		25	3						
Streptococcus pneumoniae	23		9	6	2	6				
Streptococcus faecalis	53							3	39	11
H. influenzae	98						20	41	29	8
N. gonorrhoeae	13		1	3		3	1	5		

Table II. *In Vitro* Activity of Amoxicillin Against Gram-Negative Bacilli

Organism	No. of Strains	Minimum Inhibitory Concentration ($\mu\text{g}/\text{mL}$)							
		1.25 or less	2.5	5.0	12.5	25	50	100	> 100
Proteus mirabilis	90	38	28	11					13
Shigella sonnei	26		4	11	4		1	1	5
Salmonella species	20	10	8						2
Klebsiella-Enterobacter	29		1				1	2	25
Serratia marcescens	18			1		1	3	6	7
E. coli	206	5	13	115	46	2	1	1	23

The minimum inhibitory concentrations of amoxicillin against all micro-organisms with the exception of 5 strains of *Streptococcus pneumoniae* were measured by serial dilution in agar.³⁶ The minimum inhibitory concentration against these strains of *Streptococcus pneumoniae* was estimated using the tube dilution method with Levinthal's medium.³⁴

16 NON-CLINICAL TOXICOLOGY

Acute Toxicity

The following LD₅₀ values for amoxicillin expressed in mg/kg of body weight have been reported.

Species	Route of Administration		
	P.O.	I.P.	S.C.
Mouse	> 10,000	4350	> 6,000
Rat	> 8,000	4900	> 6,000
Dog	> 3,000	-	-

Sub-acute Toxicity

Rats:

In one study male and female rats were orally administered 500 mg/kg amoxicillin daily for 21 days. With the exception of significantly greater ($p < 0.01$) BUN values in the female test group compared with controls, there were no toxic effects on the organs, tissues or fluids of the body, nor any adverse effects on food consumption, weight gain, or efficiency of food utilization reported in the study.

Histopathologic evaluation of tissues revealed a minimal degree of fatty change in livers of treated females. However, this finding was not considered a toxic change but related to a possible alteration in the intestinal flora.

Dogs:

One male and one female dog were dosed orally with 250 mg/kg amoxicillin daily for 14 days. During the period of observation, no deaths occurred, no adverse changes in body weight and no effect on food consumption was found. Laboratory values were found within normal limits. At post-mortem, no gross or microscopic abnormalities were reported and organ weights were within normal limits.

Chronic Toxicity

Rats:

In one study male and female rats were given oral doses of 200, 500 and 2000 mg/kg/day amoxicillin, 6 days a week for 26 weeks. No apparent disturbances in absolute organ weights of either treated male or female animals were noted nor was any histologic evidence of response to treatment observed.

In another study, 3 groups of Sprague-Dawley rats were given oral doses of 200, 500 and 2000 mg/kg of amoxicillin for a test period of 13-15 weeks. There were no gross or histologic changes observed in the treated rats that were considered related to the administration of amoxicillin. Some of the intermediate and low-dose groups were shown to exhibit body weight gains lower (males) or slightly higher (females) than those of the control animals.

Dogs:

It has been reported that amoxicillin was administered orally at doses of 200, 500 and 2000 mg/kg/day to male and female dogs for a period of 6 months. (Groups consisted of 6 male and 6 female dogs initially, but after 3 months dosing, each group was reduced to 3 dogs).

During the first six weeks of treatment, occasional bouts of vomiting, one to four hours after dosing, were reported in dogs receiving 2000 mg/kg/day and 4 bouts of vomiting were recorded in dogs receiving the intermediate dose of 500 mg/kg/day. Grey coloured feces were seen on very isolated occasions in dogs treated at high and intermediate dose levels only. On seven occasions it involved dogs receiving the highest dose level (2000 mg/kg/day) and on three occasions dogs receiving the intermediate dose level (500 mg/kg/day).

Body weight gains of treated males were reported to be not significantly different from those of controls, but all dosed females increased in weight at a significantly slower rate than did the controls. This factor was reported to be attributable to excessive weight gain in the control animals. Food and water consumption was not affected. No abnormalities of the eyes were observed attributable to amoxicillin.

In a second study 2 groups of Beagle dogs were given oral doses of 500 mg/kg and 200 mg/kg of amoxicillin for 13 weeks. There were no gross or histologic changes reported in the treated dogs that were considered related to the administration of amoxicillin.

Reproductive and Developmental Toxicology:

Rats:

Daily doses of 200 and 500 mg/kg amoxicillin were administered orally in one reported study. Male rats that had attained a minimum age of 40 days were treated for 63 days and sexually mature females for 14 days prior to mating. Dosing continued throughout the remainder of the investigation. The duration of gestation was unaffected by treatment at either dosage. It was noted that pregnancy rate at 500 mg/kg was slightly lower than that of controls at the first and second matings. At 200 mg/kg, the pregnancy rate was essentially comparable to control values at both matings. The chronologic sequence of mating was comparable for all groups; at 500 mg/kg the total number of animals showing evidence of mating was slightly lower than that of controls at both pairings. Pre- and post-implantation losses were comparable for all groups at the first and second pregnancies.

Among the rats allowed to rear their young, litter sizes, litter weights, mean pup weights and the pup mortality rates for the group dosed at 500 mg/kg amoxicillin were comparable to control values at birth, 4 and 21 days postpartum. Mean pup weights and pup mortality rates were similarly unaffected by 200 mg/kg amoxicillin; but litter sizes and litter weights were lower than control values from birth through lactation. These differences were considered to be unrelated to treatment. No abnormal young were observed.

Effects on Pregnancy

Mice:

It has been reported that amoxicillin administered at doses of 200, 500 and 2000 mg/kg/day orally during days 6-15 of pregnancy produced no obvious signs of reaction to treatment or deaths among parent animals. Body weight changes of pregnant dams were comparable for all groups, as was the pregnancy rate.

Fetal loss was significantly higher among all test groups than among controls. However, as implantation rates also tended to be higher at the 500 and 2000 mg/kg doses, litter sizes were only marginally, and not significantly, lower than the control value. Litter sizes and implantation rate also tended to lie at or above the upper limit of the laboratory range. Due to the latter factors, the biologic importance of the increased fetal loss was uncertain. It was noted that mean pup weights were comparable for all groups. The distribution of skeletal variants was considered to be unaffected by treatment at any dosage. A significantly higher proportion of pups with cervical ribs was found in the 200 mg/kg dose group. Cervical rib and 14th rib are the prolongations of the transverse processes of the cervical or lumbar vertebrae. Supernumerary ribs have an incidence which depends on the strain of animals. Cervical ribs are not abnormalities and have no pathologic significance.

In this experiment the incidence of cervical ribs was 12% in control rats and 16% in the drug-treated groups if the three groups are calculated together. If the groups are considered individually, then in the lowest dose group (200 mg/kg) the incidence of cervical ribs was 24%, which is, statistically, significantly higher than in the controls. This finding was not considered to be drug related since at the 500 mg/kg dose level the incidence of cervical ribs was significantly lower than in controls. At the highest dose level (2000 mg/kg) the incidence of cervical ribs was 17%, similar to the controls. The incidence of visceral abnormalities was not significantly affected at any dose level.

Rats:

Amoxicillin was administered at doses of 200 mg/kg, 500 mg/kg and 1000 mg/kg orally during gestation from day 6 through 15. Amoxicillin did not modify pregnancy, percentage of resorption and did not produce fetal abnormalities as compared with negative control rats.

Effects on Peri- and Post-Natal Development of the Rat

Amoxicillin was administered orally at 200 and 500 mg/kg/day from day 15 of gestation through lactation to 21 days post-partum. Body weight gain, pregnancy rate, and the duration of gestation of parent animals were unaffected by treatment at any dosage. There was a significant dose-related trend to lower litter size and weight at birth. This persisted through lactation to weaning despite reduced pup mortality and increased mean pup weight in the test groups compared with controls. No abnormal young were observed.

PATIENT MEDICATION INFORMATION

READ THIS FOR SAFE AND EFFECTIVE USE OF YOUR MEDICINE

Pr NOVAMOXIN

amoxicillin (as amoxicillin trihydrate)

Read this carefully before you start taking **NOVAMOXIN** and each time you get a refill. This leaflet is a summary and will not tell you everything about this drug. Talk to your healthcare professional about your medical condition and treatment and ask if there is any new information about **NOVAMOXIN**.

What is NOVAMOXIN used for?

NOVAMOXIN is used to treat certain bacterial infections. It may also be used to prevent infections in:

- Mouth, nose, tonsils and throat.
- Heart.
- Emergency situation.

How does NOVAMOXIN work?

NOVAMOXIN interferes with bacterial cell wall. This helps to:

- Stop growth of bacteria.
- Kill the bacteria.
- Reduce the infection.

Some infections are caused by viruses, such as the common cold. NOVAMOXIN **does not** kill viruses.

What are the ingredients in NOVAMOXIN?

Medicinal ingredients: amoxicillin (as amoxicillin trihydrate)

Non-medicinal ingredients:

250 mg and 500 mg capsules: Colloidal Silicon Dioxide, Dry-Flo Starch, Magnesium stearate, Sodium lauryl sulfate. Gelatine Capsules (EGC Size #2) and (EGC Size #0) : D&C Yellow # 10, FD & C Blue #1, FD & C Red #3, FD & C Red #40, FD & C Yellow # 6, Titanium Dioxide, Gelatin and White S-1-7085 (250 mg) / White SB-0007P (500 mg)

250 mg tablets: Artificial Cherry Flavor, Aspartame, Colloidal Silicon Dioxide, Dipac White (Compressible Sugar), D&C Red #30 AL Lake, FD&C Red #40 Lake, Magnesium Stearate, Mannitol, Microcrystalline Cellulose

250 mg/5 mL Granules for Oral Suspension: Artificial Cherry Raspberry Flavor, FD&C Red #40, Silicon Dioxide, Sodium Benzoate, Sodium Citrate, Sucrose, Xanthan Gum

250 mg/5 mL Sugar-Reduced Granules for Oral Suspension: Colloidal Silicon Dioxide, Mannitol 2080, Nipagin-M Sodium, Nipazol-M Sodium, Sodium Citrate, Sucrose, Xanthan Gum

For 250 mg/5 mL Sugar-Reduced Granules for Oral Suspension: D&C Yellow #10 & Artificial Banana Flavour

If you are on a special diet, or if you are allergic to any substance, ask your doctor or pharmacist whether any of these ingredients may cause a problem.

NOVAMOXIN comes in the following dosage forms:

NOVAMOXIN is available in four different forms, such as capsules, chewable tablets, Granules for Oral Suspension

and Sugar-Reduced Granules for Oral Suspension

- NOVAMOXIN 250 mg Capsules in # 2 capsules with opaque scarlet cap and yellow body
- NOVAMOXIN 500 mg Capsules in #0 capsules with opaque scarlet cap and yellow body
- NOVAMOXIN 250 mg Tablets are a rose coloured, mottled, oval shaped, single-scored compressed tablet with cherry flavour, engraved 'NOVO' with scoreline on one side and '250' on the other side
- NOVAMOXIN Granules for Oral Suspension 250 mg/5 mL
- NOVAMOXIN Sugar-Reduced Granules for Oral Suspension 250 mg/5 mL

Do not use NOVAMOXIN if:

- You have any allergies to this drug or to its ingredients (See "What are the ingredients in Novamoxin?").
- You have allergy to packaging components of this drug.
- You have allergy to penicillins, cephalosporins or similar antibiotics such as amoxicillin, ampicillin, cephalexin and others.
- You have a mononucleosis (either suspected or confirmed).

To help avoid side effects and ensure proper use, talk to your healthcare professional before you take NOVAMOXIN. Talk about any health conditions or problems you may have, including if you:

- have a history of hypersensitivity reactions to beta-lactams (ampicillin, piperacillin, etc). See "What are the possible side effects from using Novamoxin?".
- have been taken blood thinners (such as warfarin, etc.).
- have a history of mild diarrhea or colitis influenced by the use of antibiotics.
- have kidney problems.
- are pregnant or plan to become pregnant.
- are breastfeeding or plan to breastfeed. Talk to your doctor about how to feed your baby while you are taking **NOVAMOXIN**.

Tell your healthcare professional about all the medicines you take, including any drugs, vitamins, minerals, natural supplements or alternative medicines.

The following may interact with NOVAMOXIN:

- anti-cancer drug (such as Methotrexate).
- medicines used for heartburn or gout (such as probenecid, cimetidine, etc.).
- blood thinner medications (such as warfarin, etc.) that used to thin the blood and prevent clots – may predispose you to the development of bleeding problems.
- birth control pills (it may reduce effect of contraceptives).
- antibacterial medicines (such as tetracyclines) may lower effectiveness of NOVAMOXIN.

How to take NOVAMOXIN:

Antibacterial drugs like NOVAMOXIN treat only bacterial infections. They do not treat viral infections. Although you may feel better early in the treatment, NOVAMOXIN should be used exactly as directed. Misuse or overuse of NOVAMOXIN could lead to the growth of bacterial that will not be killed by NOVAMOXIN (resistance). This means that NOVAMOXIN may not work in the future.

Do not share your medicine.

Ask your pharmacist about the other products you take. Some medicines will affect the way that your body absorbs NOVAMOXIN.

Usual adult dose:

For infections: 250 mg – 500 mg every 8 hours or a single dose of 3 g.

For prevention: 3 g once before procedure, then 1.5 g every 6 hours.

Usual children’s dose:

Your doctor will tell you how much Novamoxin to give your child based on their weight and the severity of their infection. The children’s dose should not exceed the adult dose. For children over 20 kg, the adult dose should be used.

For the oral suspension, please administer using the syringe provided by your pharmacist to ensure the correct dose is given.

Take this medication by mouth as directed by your doctor.

Take Novamoxin between meals with a glass of water.

Tell your doctor if your condition does not improve.

Overdose:

If you think you or person you are caring for, have taken too much **NOVAMOXIN**, contact a healthcare professional, hospital emergency department or regional poison control centre immediately, even if there are no symptoms.

Symptoms of overdose may include: severe dizziness.

Missed Dose:

If you miss a dose, take it as soon as you remember. If it is near the time of the next dose, skip the missed dose and resume your usual dosing schedule. Do not double the dose to catch up.

What are possible side effects from using NOVAMOXIN?

These are not all the possible side effects you may feel when taking **NOVAMOXIN**. If you experience any side effects not listed here, contact your healthcare professional.

Serious side effects and what to do about them			
Symptom / effect	Talk to your healthcare professional		Stop taking drug and get immediate medical help
	Only if severe	In all cases	
COMMON			
Skin rash.			√
Skin eruption or other effect on skin or eyes.			√
Nausea.			√
Vomiting.			√
Diarrhea.			√
Bloody stool.			√
Black “hairy” tongue (glossitis).		√	
change of tooth color in children (brown, yellow or gray staining).		√	
Dizziness (light headness).		√	

Anxiety.		√	
UNCOMMON			
Hives, itch.		√	
Red rash on the face.		√	
Swelling		√	
Anaphylaxis (severe allergic reactions such as swollen nose, eyes, throat, difficulty breathing, skin blistering, rash, peeling).			√
signs of kidney problems (such as cloudy urine).			√
signs of liver problems (such as persistent nausea/vomiting, stomach/abdominal pain, unusual tiredness, yellowing eyes/skin, dark urine).			√
RARE			
Severe skin reaction (flu-like symptoms, blistering and peeling skin).			√
difficulty to fell asleep (insomnia).		√	
confusion or changes in behavior.		√	
Changes in blood cell count test results.		√	
Severe Cutaneous Adverse Reactions (SCAR) (severe skin reactions that may also affect other organs): <ul style="list-style-type: none"> • Skin peeling, scaling, or blistering (with or without pus) which may also affect your eyes, mouth, nose or genitals, itching, severe rash, bumps under the skin, skin pain, skin color changes (redness, yellowing, purplish) • Swelling and redness of eyes or face • Flu-like feeling, fever, chills, body aches, swollen glands, cough • Shortness of breath, chest pain or discomfort 			√
NOT KNOWN			
Aseptic meningitis (inflammation of the protective lining of the brain that is not caused by bacteria): confusion, fever, nausea, fatigue, sudden headache, or stiffness of your neck, sensitivity to light, vomiting			√

If you have a troublesome symptom or side effect that is not listed here or becomes bad enough to interfere with your daily activities, talk to your healthcare professional.

Reporting Side Effects

You can report any suspected side effects associated with the use of health products to Health Canada by:

- Visiting the Web page on Adverse Reaction Reporting (<https://www.canada.ca/en/health-canada/services/drugs-health-products/medeffect-canada.html>) for information on how to report online, by mail or by fax; or
- Calling toll-free at 1-866-234-2345.

NOTE: Contact your health professional if you need information about how to manage your side effects. The Canada Vigilance Program does not provide medical advice.

Storage:

For Capsules and Tablets: store between 15°C and 30°C. Blister strips should be stored between 15°C and 25°C and protected from high humidity.

Granules for oral suspension: store at room temperature between 15°C and 30°C. Keep bottle tightly closed.

Do not use after the expiry date. Generally, all expired medications should be returned to your pharmacist.

Keep out of reach and sight of children.

If you want more information about NOVAMOXIN:

- Talk to your healthcare professional
- Find the full Product Monograph that is prepared for healthcare professionals and includes this Patient Medication Information by visiting the Health Canada website (<https://www.canada.ca/en/health-canada/services/drugs-health-products/drug-products/drug-product-database.html>); the manufacturer's website <http://www.tevacanada.com>; or by calling 1-800-268-4127 ext. 3; or email druginfo@tevacanada.com.

This leaflet was prepared by Teva Canada Limited, Toronto, Ontario, M1B 2K9

Last revised: May 16, 2023